

MISOCLEAR

Misoprostol (200mcg)

High quality, effective*, proven

For early abortion of pregnancy up to 9 weeks



MARIE STOPES
INTERNATIONAL


Misoclear is high quality misoprostol.
It is recommended by doctors for early
medical abortion.

How does it work?

It causes uterine contractions and
emptying of the contents of the uterus.



How to take Misoclear Pregnancy up to 9 weeks



1 Place 4 tablets (800mcg) under the tongue (sublingually) and allow to dissolve for 30 minutes. Swallow the remainder with water

2 Place 4 tablets (800mcg) under the tongue (sublingually) and allow to dissolve for 30 minutes. Swallow the remainder with water

3 Place 4 tablets (800mcg) under the tongue (sublingually) and allow to dissolve for 30 minutes. Swallow the remainder with water

Do not take Misoclear



0 weeks → 9 weeks




10 weeks → 40 weeks

if you are more than 9 weeks pregnant



If you have severe or persistent abdominal pain



If you have an IUD in your womb

What to expect after taking Misoclear and possible side effects



Moderate bleeding



Shivering



Cramping



Vomiting/
diarrhoea

Warning signs: if experienced, seek medical help immediately These complications are rare



Heavy bleeding continues



High fever



Excess cramping/
abdominal pain



Excess vomiting/
diarrhoea

MISOCLEAR

Misoprostol (200mcg)

High quality, effective*, proven




For early abortion of pregnancy up to 9 weeks



MARIE STOPES
INTERNATIONAL

Post abortion contraception

- Speak to your doctor about contraceptive choices after an abortion.
- There is a quick return to fertility after most contraceptive methods are stopped (except for injectables and permanent methods).

When to start contraception after abortion	Method	Duration of Contraceptive Protection							Effectiveness
		1 time	1 month	3 months	3 years	5 years	10 years	Permanent	
These contraceptive methods can be started with the first pill of misoprostol	Reversible short acting contraceptives								
	 Condoms	→							85%
	 Pills	→	→						92%
	 Injectables	→	→	→					97%
	Reversible long acting contraceptives	→	→	→	→				99.9%
		→	→	→	→	→			
These contraceptive methods can be started once abortion is complete	Implants	→	→	→	→	→			
	IUD	→	→	→	→	→	→		99.2%
	Female sterilisation	→	→	→	→	→	→	→	99.5%
	Male sterilisation	→	→	→	→	→	→	→	99.8%



For information on the use of this product call:

0800 OneCall from 8am to 5pm

Or visit us at: www.misoclear.com